Contents Volume 66, 1988

Asian aridity and the zonal westerlies: Late Pleistocene and Holocene record of eolian deposition in the Northwest Pacific Ocean
D. K. Rea (Ann Arbor, MI, U.S.A.) and M. Leinen (Narragansett, RI, U.S.A.) 1
The orbital-tuned marine oxygen isotope record applied to the Middle and Late Pleistocene pollen record of Funza (Colombian Andes)
H. Hooghiemstra (Amsterdam, The Netherlands) 9
The Late Ordovician diamictic formations from southwestern Europe: north-Gondwana glaciomarine deposits
M. Robardet (Rennes, France) and F. Doré (Caen, France) 19
Reconstitution d'environnements quaternaires par l'étude de macrorestes vegetaux (pays Bateke, R.P. Congo). [Reconstruction of Quaternary environments through study of vegetable macroremains (Bateke area, P.R. Congo)]
R. Dechamps (Tervuren, Belgium), R. Lanfranchi (Libreville, Gabon), A. Le Cocq (Herbignac, France) and D. Schwartz (Pointe-Noire, R.P. Congo) 33
Fluviatile evidence for a period of Late Quaternary pluvial climate in coastal southeastern Australia
G. C. Nanson and R. W. Young (Wollongong, N.S.W., Australia) 45
Modern pollen spectra from Madagascar
D. A. Burney (Durham, NC, U.S.A.) 63
Late Pliocene to early mid-Pleistocene mammals in Eurasia: faunal succession and dispersal events
A. Azzaroli (Florence, Italy), C. De Giuli (Florence, Italy), G. Ficcarelli (Camerino, Italy) and D. Torre (Florence, Italy) 77
Palaeogeographic significance of stromatolitic buildup on Late Proterozoic platforms: the example of the West Congo Basin
R. Trompette (Marseille, France) and F. Boudzoumou (Brazzaville, Zaïre) 101
The paleoceanographic significance of the radiolarian *Didymocyrtis tetrathalamus* in eastern Cape Basin sediments
C. D. Charles and J. J. Morley (Palisades, NY, U.S.A.) 113
Fluvial processes and vertebrate taphonomy: the Upper Cretaceous Judith River Formation, south-central Dinosaur Provincial Park, Alberta, Canada
J. M. Wood, R. G. Thomas and J. Visser (Calgary, Alta., Canada) 127
Zooid size in cheilostome bryozoans as an indicator of relative palaeotemperature
B. Okamura (Halifax, N.S., Canada) and J. D. D. Bishop (London, U.K.) 145
Biotic turnover in benthic foraminifera across the Cretaceous/Tertiary boundary at El Kef, Tunisia
G. Keller (Princeton, NJ, U.S.A.) 153
Changes in the vegetation and trade winds in equatorial Northwest Africa 140,000–70,000 yr B.P. as deduced from two marine pollen records
H. Hooghiemstra (Göttingen, F.R.G.) and C. O. C. Agwu (Nsukka, Nigeria) 173
The stratigraphy, depositional environment and dating of a possible tidal surge deposit in the Beauly Firth area, northeast Scotland
B. A. Haggart (London, U.K.) 215
Ostracod shell chemistry: a new palaeoenvironmental indicator applied to a regressive/transgressive record from the Gulf of Carpentaria, Australia
P. De Deckker (Clayton, Vic., Australia), A. R. Chivas (Canberra, A.C.T., Australia), J. M. G. Shelley (Canberra, A.C.T., Australia) and T. Torgersen (Groton, CT, U.S.A.) 231
The Holocene- and Upper Pleistocene pollen record from Padul (Granada, Spain): a new study
A. Pons and M. Reille (Marseille, France) 243
Paleobiogeographic relationships of the Paleogene flora from the southeastern U.S.A.: implications for West Gondwanaland affinities
D. W. Taylor (Storrs, CT, U.S.A.) 265
Climatic implications of D/H ratios in beetle chitin
R. F. Miller (Saint John, N.B., Canada), P. Fritz (Neuherberg, F.R.G.) and A. V. Morgan (Waterloo, Ont., Canada) 277

Contents Volume 66, 1988 289